



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

ON THE RANK OF A SYMMETRICAL MATRIX.*

By L. E. DICKSON.

1. A matrix (a_{ij}) is said to be of rank r if some minor M_r of order r is not zero, while every minor M_{r+1} of order $r + 1$ is zero. A minor of order r in which there occur exactly k elements a_{ii} of the main diagonal of a square matrix (a_{ij}) shall be designated by $M_r^{(k)}$. In particular, any $M_r^{(r)}$ is called a principal minor.

Of various theorems† on the rank of a symmetrical matrix, the following theorem ‡ due to Kronecker is especially useful:

In a symmetrical matrix of rank r ($r > 0$), at least one principal minor of order r is not zero.

The following proof rests upon the fact that if n linear homogeneous equations in n variables have a set of solutions not all zero, the determinant of the coefficients is zero.

2. THEOREM. *If, in a symmetrical matrix (a_{ij}) , every $M_{r+1} = 0$ and every $M_r^{(r)} = 0$, then every $M_r = 0$.*

To proceed by induction, let ρ be a fixed integer, $0 \leq \rho < r$, for which every $M_r^{(k)} = 0$, $k > \rho$. We shall prove that every $M_r^{(\rho)} = 0$. The assumption that there is a non-vanishing $M_r^{(\rho)}$ will be shown to involve a contradiction. After a rearrangement of the rows of (a_{ij}) and the like rearrangement of the corresponding columns, a change not affecting our hypotheses, we may assume that

$$M = |a_{i1} \cdots a_{i\rho} a_{i\rho+1} \cdots a_{i2r-\rho}| \neq 0 \quad (i = 1, \dots, r).$$

Let t be one of the integers $r + 1, \dots, 2r - \rho$. Then M is a minor of

$$M_{r+1} = |a_{i1} \cdots a_{i\rho} a_{i\rho+1} a_{ir+1} \cdots a_{i2r-\rho}| = 0 \quad (i = 1, \dots, r, t).$$

Expand the latter determinant according to the elements of its last row (given by $i = t$). In the co-factor of a_{ij} ($j \geq r + 1$) appear the elements a_{kk} ($k = 1, \dots, \rho + 1$), so that these co-factors are vanishing

* Read before the American Mathematical Society, March 22, 1913.

† Cf. Bôcher's Introduction to Higher Algebra, pp. 56-59.

‡ A short proof of different nature may be found in G. Kowalewski's Determinantentheorie, p. 122-4.

$M_r^{(\rho+1)}$. Call A_j the co-factor of a_{tj} ($j \leq \rho$). Since the co-factor of $a_{t_{\rho+1}}$ is $\pm M$,

$$a_{t1}A_1 + \cdots + a_{t\rho}A_\rho \pm a_{t_{\rho+1}}M = 0 \quad (t=r+1, \dots, 2r-\rho).$$

A like equation with $t \leq r$ holds since the sum of the products of the elements of a row by the co-factors of the elements of a different row is zero. Since $M \neq 0$, it follows (end of § 1) that

$$|a_{t1} \cdots a_{t_{\rho+1}}| = 0$$

for any $\rho + 1$ values $\leq 2r - \rho$ of t . Interchange rows with columns and note that $a_{ij} = a_{ji}$. Thus

$$|a_{it}| = 0 \quad (i = 1, \dots, \rho + 1; t \text{ with any } \rho + 1 \text{ values}).$$

Hence every $M_{\rho+1}$ formed from the first $\rho + 1$ rows of M is zero, in contradiction with $M \neq 0$. Thus every $M_r^{(\rho)} = 0$ and the induction is complete.

THE UNIVERSITY OF CHICAGO,
February, 1913.